

On the scalar nonet lowest in mass

Peter Minkowski^{a *} and Wolfgang Ochs^b

^aInstitute for Theoretical Physics, University of Bern, CH-3012 Bern, Switzerland

^bMax Planck Institut für Physik, Werner Heisenberg Institut, D-80805 Munich, Germany

The hypothesis that there exists a nonet of scalars mainly composed of a valence quark-antiquark pair and mixed according to near singlet-octet separation : $f_0(980)$ singlet, $a_0^{+,0,-}(980)$, $K_0^{*+,0}(1430)$, $\bar{K}_0^{*-,-,0}(1430)$ $f_0(1500)$ octet, is put to further tests from the three body decays $D^\pm, D_s^\pm \rightarrow PS^\pm \pi^+ \pi^-$ with $PS^\pm = \pi^\pm, K^\pm$. The analysis of decay phases supports the singlet nature of $f_0(980)$.

1. Spectroscopy of $q\bar{q}$ p-waves

Following the hypothesis given in the abstract we select the following 4 p-wave nonets : $J^{PC_n} = 0^{++}, 1^{++}, 2^{++}, 1^{+-}$. They are displayed together with the pseudoscalar nonet in figure 1.




Figure 1. $J^{PC_n} = 0^{++}, 0^{+-}, 1^{++}, 2^{++}, 1^{+-}$ $q\bar{q}$ nonets.

As a consequence of the adopted selection criteria we exclude the following candidate states from the PDG listings [1]: $f_0(400 - 1200)$, $f_0(1370)$, $f_1(1410)$ and resonances with a mass exceeding $f_2(1525)$ in particular $h_1(1595)$ and $f_0(1710)$, see also [2].

We pose the following questions²:

^{*}Work supported in part by Schweizerischer Nationalfonds
²For a rather different description of low lying scalar mesons we refer e.g. to ref. [3].

- Is the flavor mixing pattern of the 0^{-+} and 0^{++} nonets similar, i.e. near singlet - octet? The two nonets would be parity doublets due to chiral symmetry, which is spontaneously and explicitly broken.

- Does the scalar gluonic meson distort through large mixing effects the scalar $q\bar{q}$ nonet beyond (spectroscopic) recognition?

In our analysis in [2] we started from the hypothesis, that the answer to this question is no. Inspecting the spectra of the p-wave nonets in Fig. 1 we observe no strong distortion in the scalar sector indeed.

- Can the last question be resolved through direct observation of the scalar gluonic meson as a conventionally narrow resonance and how reliable are the mass estimates from purely gluonic lattice-QCD [4]

$$m_{gb}(0^{++}) = 1600 \text{ MeV} \pm 10\% \quad ?$$

- Can QCD sum rules including local gluonic operators [5] shed light on the above mass estimate? The latter two questions are addressed in the accompanying paper [6].

The Gell-Mann - Okubo square mass formula

According to our analysis $f_0 \rightarrow f_<(980)$ represents the SU3 singlet, whereas

$a_0(984.7)$, $K_0(1412)$, $f_0 \rightarrow f_>(1507)$ form the associated octet.

The Gell-Mann - Okubo (first order) mass square relation then yields (in GeV^2 units)

$$\begin{aligned} m^2(f_>) &= \\ m^2(a_0) + \frac{4}{3} (m^2(K_0) - m^2(a_0)) & \quad (1) \\ 2.271 &= 0.970 + 1.365 = 2.335 \end{aligned}$$

the deviation amounts to $0.064/2.271 = 2.8\%$ ³. There is no sign - yet - of any major distortion. The degeneracy in mass of $f_<$ and a_0 , while not offending any basic principles, indicates further dynamic simplicity to be explained.

2. Further evidence for (near) octet-singlet flavor phase structure

The aim is to study and eventually confirm the nonstrange versus strange $q\bar{q}$ flavor structure of the two singlet-octet assigned isoscalar scalars :

$$\begin{aligned} f_<(980) &= \cos \vartheta | \mathbf{0} \rangle - \sin \vartheta | \mathbf{8} \rangle \\ f_>(1500) &= \sin \vartheta | \mathbf{0} \rangle + \cos \vartheta | \mathbf{8} \rangle \end{aligned}$$

The following phase convention shall be chosen in the flavor basis

$$\begin{aligned} f &= \sum_q c_q | \mathbf{q}\bar{\mathbf{q}} \rangle, \quad q = u, d, s \\ c_u &= c_d = \frac{1}{\sqrt{2}} c_{ns}; \quad c_{ns} = \sin \varphi, \quad c_s = \cos \varphi \\ | \mathbf{0} \rangle &= \frac{1}{\sqrt{3}} (1, 1, 1); \quad | \mathbf{8} \rangle = \frac{1}{\sqrt{6}} (1, 1, -2) \\ | \mathbf{ns} \rangle &= \frac{1}{\sqrt{2}} (1, 1, 0); \quad | \mathbf{s} \rangle = (0, 0, 1) \end{aligned} \quad (2)$$

and in the ns-s basis we have

$$\begin{aligned} | \mathbf{0} \rangle &= \sin \varphi_* | \mathbf{ns} \rangle + \cos \varphi_* | \mathbf{s} \rangle \\ | \mathbf{8} \rangle &= \cos \varphi_* | \mathbf{ns} \rangle - \sin \varphi_* | \mathbf{s} \rangle \\ \varphi_* &= \text{arccot} \frac{1}{\sqrt{2}} = 54.74^\circ \end{aligned} \quad (3)$$

so in the flavor basis we have

$$\begin{aligned} f_<(980) &= \sin \varphi | \mathbf{ns} \rangle + \cos \varphi | \mathbf{s} \rangle \\ f_>(1500) &= \cos \varphi | \mathbf{ns} \rangle - \sin \varphi | \mathbf{s} \rangle \\ \vartheta &= \varphi_* - \varphi \end{aligned} \quad (4)$$

As conjectured range of the singlet-octet angle ϑ we consider

$$0 \leq \vartheta \leq \vartheta_m, \quad \vartheta_m = \arcsin \frac{1}{3} = 19.47^\circ \quad (5)$$

³Eq. 1 is numerically a refinement with respect to ref. [2].

The corresponding range for φ becomes

$$35.26^\circ \leq \varphi \leq \varphi_* = 54.74^\circ \quad (6)$$

Previously [2] we have analysed the decays involving $f_0(980) : J/\Psi \rightarrow \phi f_0, \omega f_0$, the radiative decays $f_0, a_0 \rightarrow \gamma\gamma$ and $f_0 \rightarrow K\bar{K}, \pi\pi$ and concluded on a large flavor mixing similar to $\eta-\eta'$ in the pseudoscalar nonet. Now we extend our analysis to D decays.

Our analysis of the mixing pattern shall focus on the two ratios of strange to nonstrange components

$$\begin{aligned} R_< &= R(f_0(980)) = \cot \varphi \\ R_> &= R(f_0(1500)) = -\tan \varphi \end{aligned} \quad (7)$$

If the premise of mainly singlet $f_0(980)$ is correct we infer the ranges $R_< > 0$ and $R_> = -1/R_< < 0$. The relation $R_< R_> = -1$ follows from orthogonality.

The restricted ranges are

$$\begin{aligned} \frac{1}{\sqrt{2}} &\leq R_< \leq \sqrt{2} \\ -\sqrt{2} &\leq R_> \leq -\frac{1}{\sqrt{2}} \end{aligned} \quad (8)$$

We determine φ considering the three three body decays of the charmed mesons D and D_s [7]:

$$\begin{aligned} A) \quad D_s^+ &\rightarrow f_0(980) \pi^+ \\ B) \quad D^+ &\rightarrow f_0(980) \pi^+ \\ C) \quad D^+ &\rightarrow K_0^*(1430) \pi^+ \end{aligned} \quad (9)$$

We consider the color favored amplitudes ($\propto a$) which contribute to all processes in eq. 9, and color suppressed amplitudes ($\propto \epsilon a$) obtained by $\bar{d}_1 \leftrightarrow \bar{d}_2$ (see Fig. 2) and obtain

$$\begin{aligned} A &= \cos \varphi V_{ud} V_{cs}^* a \\ B &= \frac{\sin \varphi}{\sqrt{2}} V_{ud} V_{cd}^* (1 + \epsilon) a + \cos \varphi V_{us} V_{cs}^* \epsilon a \\ C &= V_{ud} V_{cs}^* (1 + \epsilon) a \end{aligned} \quad (10)$$

We consider the two ratios of partial decay widths A/B and A/C :

$$\begin{aligned} R_{A/B} &= \frac{\Gamma(D_s^+ \rightarrow f_0(980) \pi^+)}{\Gamma(D^+ \rightarrow f_0(980) \pi^+)} \\ R_{A/C} &= \frac{\Gamma(D_s^+ \rightarrow f_0(980) \pi^+)}{\Gamma(D^+ \rightarrow K_0^*(1430) \pi^+)} \end{aligned} \quad (11)$$




Figure 2. The two color suppressed quark flavor flow diagrams ($\propto \epsilon a$) in reaction B.

In the approximation $V_{ud} = V_{cs} = \cos \vartheta_c$, $V_{us} = -V_{cd} = \sin \vartheta_c$ with Cabibbo angle ϑ_c , we find

$$\begin{aligned} R_{A/B} &= 2 \frac{\Phi_1}{\Phi_2} \cot^2 \vartheta_c \cot^2 \varphi \frac{1}{|1 - (\sqrt{2} \cot \varphi - 1)\epsilon|^2} \\ R_{A/C} &= \frac{\Phi_1}{\Phi_3} \cos^2 \varphi \frac{1}{|1 + \epsilon|^2} \end{aligned} \quad (12)$$

where $\Phi_{1,2,3} = (p_{\pi^+})_{1,2,3}$ denote the phase space in s-wave decays, proportional to the π^+ momentum in the decay resonance rest frame. Using the branching fractions established by the E791 Collaboration and the PDG results we find numerically

$$\begin{aligned} \cot^2 \varphi / |1 - (\sqrt{2} \cot \varphi - 1)\epsilon|^2 &= \\ &= 1.26 \quad (1.0 \pm 0.4) \\ \cos^2 \varphi / |1 + \epsilon|^2 &= 0.52 \quad (1.0 \pm 0.3) \end{aligned} \quad (13)$$

Then we obtain as solutions two bands (a and b) due to the quadratic nature of the relations in eq.

13

$$\begin{aligned} \cot \varphi_a &= 1.11^{+0.33}_{-0.20}, \quad \varphi_a = 42.14^{+5.8}_{-7.3}^\circ \\ \epsilon_a &= (2.85 \pm 5.35) 10^{-2} \\ \tan \varphi_b &= -0.34^{+0.46}_{-0.37}, \quad \varphi_b = 161.2^{+25.7}_{-16.5}^\circ \\ \epsilon_b &= 0.31^{+0.004}_{-0.13} \end{aligned} \quad (14)$$

The angle φ is defined modulo 180° . The two solutions in eq. 14 can be distinguished through the sign of the quantity $R_<$ (or $R_>$) defined in eq. 7.

The phase (+ -) structure of $f_<$, $f_>$ is determined from interference with other resonances in D^+ and D_s^+ decays into 3π and $\pi K \bar{K}$ [7,8]. The amplitudes A, B, C in eq. 10 exhibit the + - phase structure shown in table 1.

3. Conclusions and outlook

It becomes clear from the results in table 1 and the assumed form of the amplitudes in eq. 10, that only the solution in band a) in eq. 14 is compatible with the data. This implies $\varphi = 42.14^{+5.8}_{-7.3}^\circ$ confirming the near singlet quark flavor mixing of $f_0(980)$, the ideal singlet corresponds to $\varphi_* = 54.74^\circ$.

The present analysis leaves the mixing with the scalar gluonic meson(s) completely open. Here we refer to our present results on glueballs in ref. [6]. Future work will hopefully establish the full structure of the scalar nonet lowest in mass including the scalar glueball $gb(0^{++})$.

Acknowledgement

It is a pleasure to thank all organizers and participants of the QCD2002 conference, in particular Stephan Narison, for enabling the meeting to evolve in a lively and fruitful atmosphere.

REFERENCES

- [1] K. Hagiwara et al., Phys. Rev. D66 (2002) 010001.
- [2] P. Minkowski and W. Ochs, EPJ C9 (1999) 283.
- [3] F. Close and N. Tornqvist, hep-ph/0204205.
- [4] P. Rüfenacht, hep-lat/0105033,

Table 1

Decay phases (in degrees) for resonances in D and D_s decays as measured by experiments E687 and E791. In each line an overall phase has been fixed arbitrarily. The states marked by (*) are not directly evident in the plots. Comparison with theoretical expectations for the mixing angle φ and predictions for $f_0(980)$ near flavour singlet and $f_0(1500)$ as octet partner ($0^\circ < \varphi < 90^\circ$) with arbitrary angle α , standard choice is $\alpha = 0$.

$D \rightarrow 3\pi$	$d\bar{d} \rightarrow$		
	$\rho(770)$	$f_0(980)$	$f_2(1270)^*$
E687	$27 \pm 14 \pm 11$	$197 \pm 28 \pm 24$	$207 \pm 17 \pm 4$
E791	$0(\text{fixed})$	151.8 ± 16.0	102.6 ± 16.0
Theory	$-d\bar{d}$	$d\bar{d} \sin(\varphi)$	$d\bar{d}$
	$\alpha + 180^\circ$	α	α
$D_s \rightarrow 3\pi$	$s\bar{s} \rightarrow$		
	$f_0(980)$	$f_2(1270)^*$	$f_0(1500)$
E687	$0(\text{fixed})$	83 ± 16	210 ± 10
E791	$0(\text{fixed})$	$133 \pm 13 \pm 28$	$198 \pm 19 \pm 27$
Theory	$s\bar{s} \cos(\varphi)$	$\varepsilon s\bar{s}$	$-s\bar{s} \sin(\varphi)$
	α	α	$\alpha + 180^\circ$
$D_s \rightarrow \pi K \bar{K}$	$s\bar{s} \rightarrow$		
	$f_0(980)^*$	$\phi(1020)$	
E687	$159 \pm 22 \pm 16$	$178 \pm 20 \pm 24$	
Theory	$s\bar{s} \cos(\varphi)$	$s\bar{s}$	
	α	α	

G.S. Bali, hep-ph/0110254.

- [5] S.Narison, these proceedings, hep-ph/0208081,
H.G. Dosch and S. Narison, hep-ph/0208271.
- [6] W. Ochs and P. Minkowski, these proceedings.
- [7] E.M. Aitala et al., E791 collaboration, Phys. Rev. Lett. 86, (2001) 770, hep-ex 0007028, and Phys. Rev. Lett. 86, (2001) 765, hep-ex 0007027.
- [8] P.L. Frabetti et al., E687 collaboration, Phys. Lett. B407 (1997) 79, and Phys. Lett. B351 (1995) 591.